

Web Images Maps News Shopping Gmail more ▾

Sign in



((select) (pathogenic e.coli) (acidic medium))

Search

Advanced Search
Preferences

Web Results 1 - 10 of about 23,900 for ((select) (pathogenic e.coli) (acidic medium)). (0.24 seconds)

Isolation and Molecular Characterization of Nalidixic Acid ...

Fluoroquinolone use in poultry production may select for resistant Escherichia ... From 55 products that yielded both Nal-resistant and susceptible E. coli, ... satisfied criteria for extraintestinal pathogenic E. coli (ExPEC), i.e., ...

aac.asm.org/cgi/content/abstract/47/7/2161 - Similar pages

by JF Johnson - 2003 - Cited by 47 - Related articles - All 7 versions

Isolation and Molecular Characterization of Nalidixic Acid ...

Fluoroquinolone use in poultry production may select for resistant Of the 169 chicken samples cultured, 150 (89%) yielded E. coli on nonselective media. There is ample precedent for cross-species pathogenicity of E. coli. ...

www.pubmedcentral.nih.gov/articlerender.fcgi?artid=161843 - Similar pages

by JF Johnson - 2003 - Cited by 47 - Related articles - All 7 versions

BiomedExperts: Isolation and molecular characterization of ...

'Isolation and molecular characterization of nalidixic acid-resistant extraintestinal pathogenic Escherichia coli from retail chicken products. ... Fluoroquinolone use in poultry production may select for resistant Escherichia coli that ... satisfied criteria for extraintestinal pathogenic E. coli (ExPEC), i.e., ...

www.biomedexperts.com/.../isolation_and_molecular_characterization_of_nalidixic_acid-resistant_extraintestinal_pathogenic... - 20k - Cached - Similar pages

Acid resistance of Escherichia coli O157:H7 - a review ...

then select price, enter quantity and click add to order ... evidence for acid resistance in E.coli, by first providing an overview of pathogenic E.coli and ...

www.campden.co.uk/publ/pubfiles/r28.htm - 6k - Cached - Similar pages

Pre-harvest factors influencing the acid resistance of Escherichia ...

The media was autoclaved at 121 °C for 15 min. The inoculating E. coli O157:H7 that acid resistance for both pathogenic and nonpathogenic E. coli could be Preliminary foodnet data on the incidence of foodborne illnesses—Select ...

jas.fass.org/cgi/content/full/81/4/1080 - Similar pages

by C J Fu - 2003 - Cited by 6 - Related articles - All 6 versions

Strain of Escherichia coli by bacterial conjugation - Patent 4440748

(b) inoculating a nutrient medium containing raffinose as the only carbon ... P155 Escherichia coli K88 + , sensitive to nalidixic acid, has genes responsible for ... and nalidixic acid (O-raf+nal) to select for nalidixic acid resistant cells which ... Pathogenicity of E. Coli ATCC No. 39076 For Calves and Piglets ...

www.freepatentsonline.com/4440748.html - Similar pages

by AC Graham - 1984 - Cited by 2 - Related articles - All 2 versions

Food Control : Antibacterial activity of lactic acid bacteria ...

All of which can antagonize the growth of some spoilage and pathogenic bacteria ... L. innocua and E. coli) in order to simulate the workshop ecological niche. ... and finally to select interesting strains for the role of bio-preservatives. To restrict the extent of acid production, low-glucose medium, i.e., ...

linkinghub.elsevier.com/retrieve/pii/S0956713505000551 - Similar pages

by S Ammor - 2006 - Cited by 13 - Related articles - All 5 versions

[PDF] (Microsoft Word - Competitive Exclusion of Antibiotic Resistant ...

File Format: PDF/Adobe Acrobat - [View as HTML](#)

flumequine and oxolinic acid. • A set of naive birds was given direct ... broth plus enrofloxacin to select for the antibiotic resistant bacteria. Results ... Multi-resistant Pathogenic E. coli 078:K80 (instant Seeders). ...
www.safe-poultry.com/documents/aviguard/C-E_Antibiotic_ResistantE_coli.pdf - [Similar pages](#)

[doc] [Sally Tobler](#)

File Format: Microsoft Word - [View as HTML](#)

Some strains of the E. coli are pathogenic, such as E. coli 0157:H7; ... The first test performed is the presumptive test; these tests select for any ... All three of these tests indicate coliform presence if acid and gas is produced in each. Adding methyl red to the medium will cause the medium to change to a ...
filebox.vt.edu/users/sktobler/EDCI%20portfolio/discussion6.doc - [Similar pages](#)

[Escherichia coli Serotype O157:H7: Novel Vehicles of Infection and ...](#)

Jan 18, 1996 ... The implication of acidic foods as vehicles of infection has dispelled the Though intended solely to select for serotype O157:H7, sorbitol-containing media may also exclude the isolation of other pathogenic E. coli ...
www.cdc.gov/ncidod/EID/vol1no2/feng.htm - 37k - [Cached](#) - [Similar pages](#)
by P Feng - [Cited by 108](#) - [Related articles](#) - [All 6 versions](#)

[1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

[Search within results](#) | [Language Tools](#) | [Search Tips](#) | [Dissatisfied? Help us improve](#) |
[Try Google Experimental](#)

[Google Home](#) - [Advertising Programs](#) - [Business Solutions](#) - [Privacy](#) - [About Google](#)